

Follicle Stimulating Hormone

Page 1 of 2

Diagnostic Test Information Sheet 04

Test Name: Follicle Stimulating Hormone

Test code: hFSH

Special Precautions: None

Specimen Handling: 1. Collect specimen in a 5 mL SST Gold top tube.

2. Gently invert tube 5-6 times.

3. Label specimen with:

o full name of patient

o one other unique patient identifier (i.e. DOB and/or

Medical Health Number)

o test name

4. Allow to clot upright for a minimum of 30 minutes.

5. Centrifuge for 10 minutes at 1300 RCF.

6. Place specimen in specimen transport bag.

7. Seal bag.

8. Fold requisition and place in sleeve of bag.

Reference Range: Male (>16 years old): 1.55-9.74 IU/L

Female (>16 years old):

Follicular phase: 1.98-11.6 IU/L Mid cycle: 5.14-23.4 IU/L Luteal phase: 1.38-9.58 IU/L Post-menopausal: 21.5-131 IU/L

Please note new reference ranges effective Dec 21, 2021

Turnaround Time: Not available on a STAT basis.

Routine: 72 hours

Sample Type: Serum

Sample Container: Gold Top Serum Separator Tube

Minimum Volume: 1 ml

Collection and Shipping: Refrigerate @ 2-8° C. If sent out to referral lab, send on ice pack.

Method: Beckman Coulter Access 2- Immunoenzymatic Assay

Stability Time: 2 days @ 2-8°C.

Interferences: Hemolysis: no significant interference

Lipemia: no significant interference Bilirubin: no significant interference

Revision Number:	1.0	Effective Date:	Dec 22, 2021
1			



Follicle Stimulating Hormone

Page 2 of 2

Diagnostic Test Information Sheet 04

Measurement Uncertainty: Available upon request. Please call 604-336-2090 or email

in fo@novavital aboratory.com.